Student Medical Exemption Certificate for Required Immunizations

Name of Primary Care Provider gran	ting exemption:	
Please check one (practitioner grant	ing exemption mus	t be licensed as one of the following):
\square Physician (MD or DO) \square P	hysician Assistant	☐ APRN
CT License number:		
NPI:		
Phone number:		Email:
Directions:		
Part 1. Please complete the demogr	aphics section on tl	ne patient/student.
Part 2. Please mark the contraindica apply).	tions/precautions t	that apply to this patient/student (indicate all that
Part 3 . If no contraindications or prepatient/student requires the exemp		art 2, write a brief explanation of the reason the
Part 4. Sign the Statement of Clinica	l Opinion and date	the form.
Attach a copy of the patient/studen	t's most current im	munization record.
Part 1. Patient/Student Information	:	
First name (in full)	Middle initial	Last name
Date of Birth		
Mailing Address	City	
State	Zip	
Parent/Guardian: First Name		Last name
Primary phone number		
School name		
School address		
City		
State	Zip	
Current or Grade student is entering		

Part 2. Please mark the vaccine(s), exemption duration, and all contraindications/precautions that apply to this patient/student for each vaccine.

Medical contraindications and precautions for immunizations are based upon the Advisory Committee on Immunization Practices (ACIP) <u>Comprehensive General Recommendations and Guidelines</u>, published by the Centers for Disease Control and Prevention.

A **contraindication** is a condition in a recipient that increases the risk for a serious vaccine adverse event (VAE) or compromises the ability of the vaccine to produce immunity.

A **precaution** is a condition in a recipient that might increase the risk for a serious VAE or that might compromise the ability of the vaccine to produce immunity. Under normal conditions, vaccinations are deferred when a precaution is self-limiting, but can be administered if the precaution condition improves.

CDC Recognized Contraindications and Precautions

Vaccine	Exemption Duration	ACIP Contraindications and Precautions (Check all that apply)
☐ Diphtheria-	☐ Temporary	Contraindications
Tetanus-and acellular	through:	☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
Pertussis (DTaP)	mm/ yyyy ☐ Permanent	☐ Encephalopathy within seven days after receipt of previous dose of DTP or DTaP
		Precautions
		☐ Progressive neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy: defer DTaP until neurologic status clarified and stabilized
		☐ GBS <6 weeks after previous dose of tetanus-toxoid—containing vaccine
		☐ Fever greater than 40.5°C (104.9°F) <48 hours after vaccination of previous dose of DTP or DTaP
		☐ History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid—containing or tetanus-toxoid—containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid—containing vaccine
		☐ Moderate or acute illness with or without fever
☐ Hepatitis A	☐ Temporary	Contraindications
	through:/ mm/ yyyy □ Permanent	☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose
		or to a vaccine component
		Precautions
		☐ Moderate or severe acute illness with or without fever

☐ Hepatitis B	\square Temporary	Contraindications
	through: /	☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	mm/ yyyy	☐ Hypersensitivity to yeast
	☐ Permanent	Precautions
		☐ Moderate or severe acute illness with or without fever
☐ Haemophilus influenzae type b (HiB)	☐ Temporary through:/mm/ yyyy	 Contraindications □ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component □ Age <6 weeks
	□ Permanent	Precautions
		☐ Moderate or severe acute illness with or without fever
☐ Inactivated	☐ Temporary	Contraindications
Influenza Virus (IIV)	through: /	 Severe allergic reaction (e.g., anaphylaxis) after previous dose of influenza vaccine or to vaccine component
	mm/ yyyy	Precautions
	☐ Permanent	☐ GBS <6 weeks after a previous dose of influenza vaccine
		☐ Moderate or severe acute illness with or without fever
		☐ Egg allergy other than hives, e.g., angioedema, respiratory distress, lightheadedness, recurrent emesis; or required epinephrine or another emergency medical intervention (IIV may be administered in an inpatient or outpatient medical setting and under the supervision of a health care provider who is able to recognize and manage severe allergic conditions).
☐ Inactivated	☐ Temporary	Contraindications
Polio Vaccine (IPV)	through: /	☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
	mm/ yyyy	Precautions
	☐ Permanent	☐ Pregnancy
		☐ Moderate or acute illness with or without fever

☐ Live	☐ Temporary through:	Contraindications			
Attenuated Influenza Virus (LAIV)		☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component			
(LAIV)	mm/ yyyy	☐ Concomitant use of aspirin or aspirin-containing medication in children and adolescents			
		☐ LAIV4 should not be administered to persons who have taken oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days.(e)			
		☐ Pregnancy			
		☐ Children aged 2 through 4 years who have received a diagnosis of asthma or whose parents or caregivers report that a health care provider has told them during the preceding 12 months that their child had wheezing or asthma or whose medical record indicates a wheezing episode has occurred during the preceding 12 months.			
		☐ Persons with active cerebrospinal fluid/oropharyngeal communications/leaks.			
		☐ Close contacts and caregivers of severely immunosuppressed persons who require a protected environment.			
		☐ Persons with cochlear implants (due to the potential for CSF leak, which might exist for some period of time after implantation. Providers might consider consultation with a specialist concerning risk of persistent CSF leak if an age-appropriate inactivated or recombinant vaccine cannot be used).			
		☐ Altered Immunocompetence			
		☐ Anatomic or functional asplenia (e.g. sickle cell disease)			
		Precautions			
		☐ GBS <6 weeks after a previous dose of influenza vaccine			
		☐ Asthma in persons aged 5 years old or older			
		☐ Medical conditions which might predispose to higher risk of complications attributable to influenza(d)			
		☐ Moderate or severe acute illness with or without fever			
☐ Meningo-	☐ Temporary	Contraindications			
coccal conjugate vaccines		☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component, including yeast			
(MenACWY)		Precautions			
		☐ Moderate or severe acute illness with or without fever			

☐ Measles-	☐ Temporary through:	Contraindications				
Mumps-Rubella (MMR)		☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or				
	mm/ yyyy	to a vaccine component Pregnancy				
	☐ Permanent					
		☐ Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised)				
		☐ Family history of altered immunocompetence (i)				
		Precautions				
		☐ Recent (≤11 months) receipt of antibody-containing blood product (specific interval depends on product)				
		☐ History of thrombocytopenia or thrombocytopenic purpura				
		☐ Need for tuberculin skin testing or interferon-gamma release assay (IGRA) testing (k)				
		☐ Moderate or severe acute illness with or without fever				
☐ Pneumo-	☐ Temporary	Contraindications				
coccal (PCV13)	through:/ mm/ yyyy	☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose of PCV13 or any diphtheria-toxoid—containing vaccine or to a component of a vaccine (PCV13 or any diphtheria-toxoid—containing vaccine), including yeast				
	☐ Permanent	Precautions				
		☐ Moderate or acute illness with or without fever				
☐ Tdap	☐ Temporary	Contraindications				
raup	through:	☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component				
	mm/ yyyy ☐ Permanent	☐ Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, or Tdap				
		Precautions				
		☐ GBS <6 weeks after a previous dose of tetanus-toxoid—containing vaccine				
	l	Drogressive or unstable neurological disorder, uncontrolled				
		☐ Progressive or unstable neurological disorder, uncontrolled seizures, or progressive encephalopathy until a treatment regimen has been established and the condition has stabilized				

		dose of diphtheria-toxoid—containing or tetanus-toxoid—containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid—containing vaccine			
		☐ Moderate or severe acute illness with or without fever			
☐ Varicella ☐ Temporary		Contraindications			
	through:	☐ Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component			
mm/ yyyy		☐ Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy (i) or patients with HIV infection who are severely immunocompromised) (g)			
		☐ Pregnancy			
		☐ Family history of altered immunocompetence (j)			
		Precautions			
		☐ Recent (<11 months) receipt of antibody-containing blood product (specific interval depends on product)			
		☐ Moderate or acute illness with or without fever			
=	a for a contraindic	nedical exemption for a vaccine based on a condition that does meet any ration or precaution listed in part 2.			
	•	t the allergic or other reaction for which medical exemption is being he following that apply:			
\square This patient h	as an autoimmun	e disorder			
\square This patient h	as a family history	y of an autoimmune disorder			
\square This patient h	as a family history	of a reaction to a vaccination			
☐ This patient h testing	as a genetic predi	sposition to a reaction to a vaccination as determined through genetic			
\square This patient h	as a previous doci	umented reaction that is correlated to a vaccination			
☐ Other conditi	on/reaction not lis	sted above (must specify):			
Please provide ar					

Part 4. Statement of Clinical Opinion

In accord with the	legal requirements of	Public Act 21-6,	the vaccine(s)	indicated at	oove is/are	e in my cli	nical
opinion medically of	contraindicated for th	is patient/studen	nt due to the ph	nysical condi	ition as ex	plained at	oove.

Clinician's Signature			
_			
Date			

A person may be placed into quarantine or isolation when there are "reasonable grounds to believe [a person] to be infected with, or exposed to, a communicable disease or to be contaminated or exposed to contamination or at reasonable risk of having a communicable disease or being contaminated or passing such communicable disease or contamination to other persons if the commissioner determines that such individual or individuals pose a significant threat to the public health and that quarantine or isolation is necessary and the least restrictive alternative to protect or preserve the public health." Conn. Gen. Stat. § 19a-131b(a).